Astronomy & Astrophysics manuscript no. ngc6388˙ref
September 28, 2011

© ESO 2011

arXiv:1109.5719v1 [astro-ph.HE] 26 Sep 2011

Letter to the Editor

IGRJ17361-4441: a possible new accreting X-ray binary in NGC6388

E. Bozzo[1], C. Ferrigno[1], J. Stevens[2], T. M. Belloni[3], J. Rodriguez[4], P. R. den Hartog[5], A. Papitto[6], I. Kreykenbohm[7], F. Fontani[8], and L. Gibaud[1]

[1] ISDC Data Centre for Astrophysics, Chemin dEcogia 16, CH-1290 Versoix, Switzerland; e-mail: `enrico.bozzo@unige.ch`
[2] CSIRO Astronomy & Space Science, Australia Telescope National Facility, P.O. Box 76, Epping, NSW 1710, Australia
[3] INAF, Osservatorio Astronomico di Brera, via E. Bianchi 46, I-23807 Merate (LC), Italy
[4] Laboratoire AIM, CEA/IRFU - CNRS/INSU - Université Paris Diderot, CEA DSM/IRFU/SAp, F-91191 Gif-sur-Yvette, France
[5] Stanford University HEPL/KIPAC Physics, 382 Via Pueblo Mall Stanford, 94305, USA
[6] Dipartimento di Fisica, Universitá degli Studi di Cagliari, SP Monserrato-Sestu, KM 0.7, 09042 Monserrato, Italy
[7] Dr. Karl Remeis-Sternwarte & Erlangen Centre for Astroparticel Physics, Sternwartstr. 7, 96049 Bamberg
[8] INAF - Observatorio Astrofisico di Arcetri, L.go E. Fermi 5, I-50125, Firenze, Italy

Submitted: 2011 September 5 - Accepted: 2011 September 22

ABSTRACT

IGR J17361-4441 is a newly discovered *INTEGRAL* hard X-ray transient, located in the globular cluster NGC6388. We report here the results of the X-ray and radio observations performed with *Swift*, *INTEGRAL*, *RXTE*, and the Australia Telescope Compact Array (ATCA) after the discovery of the source on 2011 August 11. In the X-ray domain, IGR J17361-4441 showed virtually constant flux and spectral parameters up to 18 days from the onset of the outburst. The broad-band (0.5-100 keV) spectrum of the source could be reasonably well described by using an absorbed power-law component with a high energy cut-off ($N_{\rm H}$≃0.8×10^{22} cm^{-2}, Γ≃0.7-1.0, and $E_{\rm cut}$≃25 keV) and displayed some evidence of a soft component below ~2 keV. No coherent timing features were found in the *RXTE* data. The ATCA observation did not detect significant radio emission from IGR J17361-4441, and provided the most stringent upper limit (rms 14.1 μJy at 5.5 GHz) to date on the presence of any radio source close to the NGC6388 center of gravity. The improved position of IGR J17361-4441 in outburst determined from a recent target of opportunity observation with *Chandra*, together with the X-ray flux and radio upper limits measured in the direction of the source, argue against its association with the putative intermediate-mass black hole residing in the globular cluster and with the general hypothesis that the *INTEGRAL* source is a black hole candidate. IGR J17361-4441 might be more likely a new X-ray binary hosting an accreting neutron star. The ATCA radio non-detection also permits us to derive an upper limit to the mass of the suspected intermediate massive black hole in NGC6388 of $\lesssim$600 $M_{\odot}$. This is a factor of 2.5 lower than the limit reported previously.

Key words. gamma rays: observations – X-rays: individuals: IGR J17361-4441

1. Introduction

IGR J17361-4441 was discovered by the IBIS/ISGRI telescope (Ubertini et al. 2003) onboard *INTEGRAL* (Winkler et al. 2003) on 2011 August 11 (Gibaud et al. 2011). The source was located in the direction of the globular cluster (GC) NGC6388 and detected at an X-ray flux of 9.7×10^{-11} erg cm^{-2} s^{-1} (20-100 keV). Follow-up observations carried out with *Swift* and *Chandra* (Ferrigno et al. 2011b; Pooley et al. 2011) provided a first description of the source X-ray emission in the soft energy band (0.3-10 keV), a refined source position at RA = 17:36:17.418, Dec = -44:44:05.98 (J2000; the nominal *Chandra* positional accuracy is 0.6 arcsec, 90% c.l.) and confirmed its localization in NGC6388. In the past, several X-ray observational campaigns have been carried out in the direction of this GC, as it is believed to host an intermediate-mass black hole (IMBH) at its center of gravity (hereafter COG; Lanzoni et al. 2007). Here we report on the analysis of all the available X-ray data collected during the first 18 days of the outburst observed from IGR J17361-4441, and use a simultaneous observation in the radio domain with the ATCA telescope to investigate the nature of the source.

2. Data analysis

2.1. INTEGRAL

We analyzed all the available data for the IBIS/ISGRI (20-150 keV, Lebrun et al. 2003) and for the JEM-X telescope (5-20 keV, Lund et al. 2003) onboard *INTEGRAL* that were performed in the direction of IGR J17361-4441 from 2011 August 11 (rev. 1078) to 2011 August 29 (rev. 1083). Data analysis was carried out by using version 9.0 of the OSA software distributed by the ISDC (Courvoisier et al. 2003). In Fig. 1, we show the IBIS/ISGRI mosaicked significance map around IGR J17361-4441 in the 20-100 keV energy band (total exposure time ~500 ks). In this mosaic, IGR J17361-4441 is detected at a significance level of 15 σ at an average flux of 1.2×10^{-10} erg/cm^2/s. The corresponding IBIS/ISGRI spectrum is discussed later in Sect. 2.3. From rev. 1078 to 1083, IGR J17361-4441 was inside the field of view (FOV) of the X-ray monitor onboard *INTEGRAL*, JEM-X, only for a total effective exposure time of 17 ks and was not detected. We derived a 3 σ upper limit to the 3-35 keV X-ray flux from IGR J17361-4441 of 3.8 mCrab[1] (corresponding to 1.1×10^{-10} erg/cm^2/s). This is compatible with the

---

[1] Calculated by using the calibration observation of the Crab in satellite rev. 967 (from 2010 September 13 to 15)



Fig. 1. IBIS/ISGRI FOV around IGR J17361-4441 (20-100 keV energy band). We used data from rev. 1078 to 1083. The horizontal band on the bottom of the image shows the colour-coded significances in the mosaic expressed in units of σ.

value expected based on the broad-band spectral properties of the source (see Sect. 2.3).

2.2. RXTE

A follow-up observation of IGR J17361-4441 was performed with the PCA onboard *RXTE* (Bradt et al. 1993) on 2011 August 17 (total exposure time 1.6 ks). The pointing was carried out with an offset of ~0.53 degrees from the nominal position of IGR J17361-4441 to limit[2] the contamination from the bright nearby low mass X-ray binary 4U 1735−444. The good time intervals were based on a maximum offset from the source of 0.55 degrees, a minimum elevation of ten degrees over the Earth limb, an electron ratio lower than 0.2, and at least ten minutes having passed since the SAA passage (see also, e.g. Ferrigno et al. 2011a). We analyzed the PCA event mode data in the GoodXenon configuration, which has a time resolution of 0.96μs and selected all events from the PCU 1, 2, and 4 (PCU0 suffered a breakdown during the observation). The event arrival times were all barycenter-corrected to the Solar System center-of-mass using the available tool FAXBARY and the most-accurately determined source position in *Chandra* data (see Sect. 1). We constructed a power density spectrum after accumulating light curves in bins of 0.49 ms, so that frequencies of up to ~1 kHz could be explored when searching for coherent signals (see Fig. 2). To maximize the frequency resolution, we used a single interval to compute the fast Fourier transform (FFT). We did not find any significant peak at a 3σ c.l. (we followed the approach described in Vaughan et al. 1994). The high background level of the PCA observation and the contamination from 4U 1735−444 did not permit an accurate estimation of the corresponding pulsed fraction upper limit.



Fig. 2. Power spectral density obtained from the PCA events of PCU 1,2, and 4 (we used a single time interval with bins of 0.49 ms). The dashed line represents the 3σ limit for detection of coherent signals.

2.3. Swift

Swift/XRT monitored the outburst of IGR J17361-4441 starting from 2011 August 16 for about 18 days (the last observation considered here was carried out on 2011 September 2 at 06:10 UTC). We processed all the *Swift*/XRT data with standard procedures (Burrows et al. 2005) and the latest calibration files available (caldb v. 20110725). All observations were performed in photon-counting mode (PC, time resolution 2.5 s). The data analysis technique that we used is the same as that described in Bozzo et al. (2009). All the XRT spectra extracted from the available observations could be well accurately reproduced by using an absorbed power-law model with $N_{\rm H}\simeq(0.2\text{-}0.5)\times10^{22}$ cm^{-2} and $\Gamma\sim0.5\text{-}0.9$. During this period, the source displayed a virtually constant X-ray flux of $(4.5\text{-}4.8)\times10^{-11}$ erg/cm^2/s (1-10 keV not corrected for absorption). The XRT lightcurve of the outburst is reported in Fig. 3. By using the *Swift*/XRT on-line analysis tool[3] with all the XRT data available in the direction of NGC6388, we also determined the new XRT position of IGR J17361-4441 to be RA=17:36:17.27, Dec=-44:44:07.0, with an associated uncertainty of 1.9 arcsec (90% c.l.; see Fig. 5). This is consistent with the position reported previously by Ferrigno et al. (2011b) and the refined *Chandra* position reported by Pooley et al. (2011). We studied the broad-band spectral properties of the X-ray emission from IGR J17361-4441 by combining XRT data carried out before 2011 August 29 with those obtained during the same period with ISGRI (rev. 1078 to 1083). The joint XRT+ISGRI spectrum is shown in Fig. 4. We fit the two spectra together by using an absorbed power-law (PL) model with a cut-off at high energy (CUTOFFPL in XSPEC). A blackbody component (BB) was required to take into account the residuals at the lower energies ($\lesssim$2 keV). A normalization constant (fixed to 1 for the XRT spectrum) was included in the fit to account for the intercalibration between the different instruments and the variability (if any) of the source. From the fit ($\chi^2_{\rm red}$/d.o.f.=0.95/269), we estimated that $N_{\rm H}$=0.8±0.1 erg/cm^2/s, Γ=1.0±0.1, $E_{\rm cut}=24^{+20}_{-9}$ keV, $kT_{\rm BB}$=0.076±0.005 keV, and $R_{\rm BB}=1.4^{+0.8}_{-0.5}\times10^3$ $d_{10\ kpc}$ km (where $E_{\rm cut}$ is the cut-off energy, $d_{10\ kpc}$ is the source distance in units of 10 kpc, and all the uncertainties are given at the 90% c.l.). The normalization constant used for ISGRI turned out to be

---

[2] By using archival PCA observations, we estimated an average on-axis count-rate over three active PCUs of ~ 1400 cts/s for 4U 1735−444. Therefore, this source might contribute up to ~25% of the signal from the observation of IGR J17361-4441, which has a background-subtracted count-rate of ~ 10 cts/s.

[3] See http://www.swift.ac.uk/user_objects/; see also Evans et al. (2009).



Fig. 3. Lightcurve of IGR J17361-4441 as observed by *Swift*/XRT (0.3-10 keV). The bin time is 250 s and the start time is 2011 August 16 at 17:01 (UTC).



Fig. 4. Broad-band spectrum of IGR J17361-4441 obtained by using data from *Swift*/XRT and *INTEGRAL*/ISGRI. The best-fit model (absorbed cut-off PL plus a BB component, see text for details) and the residuals from this fit are also shown.

$C=1.2^{+0.8}_{-0.5}$ and no significant changes in the spectral fit parameters were observed by fixing $C=1$. The source average X-ray flux was $(4.6^{+0.1}_{-0.5})\times10^{-11}$ erg/cm^2/s in the 1-10 keV energy band and $7.8^{+0.8}_{-3.8}\times10^{-11}$ erg/cm^2/s in the 20-100 keV energy band. Compatible results were obtained by fitting the *RXTE*/PCA spectrum with the model described above. The 3-15 keV X-ray flux measured from the PCA was $6.7^{+0.1}_{-3.4}\times10^{-11}$ erg/cm^2/s, i.e. slightly higher than that extrapolated from the XRT energy band. Part of this discrepancy might be due to the limited time range spanned by the PCA data with respect to XRT and ISGRI, and possible contamination of the nearby bright persistent low mass X-ray binary 4U 1735-444 (located ∼0.5 deg away from IGR J17361-4441, see also Sect. 2.2).

2.4. ATCA

To hel us establish the nature of IGR J17361-4441, we also obtained on 2011 August 25 a 8 h-long target of opportunity observation (ToO) with the Australia Telescope Compact Array (ATCA; Wilson et al. 2011) in configuration 6D (baseline is 6 km). The observation was conducted with a 2 GHz bandwidth centered around 5.5 GHz, 9 GHz, 17 GHz, and 19 GHz. The total integration time was about 5.3 h for the two lower frequencies (observed simultaneously) and 2.7 h for the higher frequencies (observed simultaneously). Natural-weighted images were made from the ATCA data, with FOVs (FWHM) of 8.5, 5.2, 2.8, and 2.5 arcmin, and synthesized beam sizes (FWHM) of 4.5×1.8, 2.7×1.1, 1.7×0.6, and 1.5×0.5 arcsec at 5.5, 9, 17, and 19 GHz, respectively. No significant radio emission was found around the X-ray position of IGR J17361-4441. The rms noise levels of the images were 14.1 μJy, 19.0 μJy, 20.0 μJy, and 21.2 μJy for the 5.5 GHz, 9 GHz, 17 GHz, and 19 GHz frequencies, respectively. The rms noise level corresponding to the image in the 5.5 Ghz frequency is a factor of ∼2.5 lower than that reported previously by Cseh et al. (2010, see also Sect. 3).



Fig. 5. Close view of the *Chandra* archival image (OBSID. 5055; 0.5-10 keV) around the core of NGC6388. In the image, black circles correspond to all the *Chandra* sources detected so far in quiescence by Cseh et al. (position uncertainty 0.5 arcsec; 2010). The blue circle represents the COG of the GC (position uncertainty 0.3 arcsec). The improved XRT position determined in the present work is indicated with a red circle (see Sect. 2.3), and the *Chandra* position reported by Pooley et al. (2011) is represented with a green circle.

3. Discussion

The X-ray monitoring campaign triggered after the discovery of IGR J17361-4441 with *INTEGRAL*, has enabled us to significantly improve the accuracy of the source position, and confirm its location within the core of the GC NGC6388 (the COG of the GC has coordinates RA=17:36:17.23, Dec=-44:44:07.1; the estimated uncertainty is 0.3 arcsec; Lanzoni et al. 2007, see also Sect. 2.3). The XRT data have helped us to demonstrate that the source remained at a virtually constant flux during the first 18 days after the onset of the outburst, and did not display any significant spectral variability. We have been able to fit the broad-band (0.5-100 keV) spectrum of the source reasonably well using a cut-off PL model ($N_{\rm H}\simeq0.8\times10^{22}$ cm^{-2}, $\Gamma\simeq0.7$-1.0, and $E_{\rm cut}\simeq25$ keV) and displayed some hint of a soft spectral component below ∼2 keV. The estimated flux in the 1-100 keV energy band is 1.7×10^{-10} erg/cm^2/s, which corresponds to a luminosity of $L_{\rm X}\simeq3.6\times10^{36}$ erg/s at the distance of NGC6388 (13.2 kpc; Dalessandro et al. 2008).

The GC NGC6388 was previously observed with *Swift*, *XMM–Newton*, and *Chandra*, because it is believed to host an IMBH in its COG (see also, Nucita et al. 2008, and references therein). To date, the deepest X-ray observations of the GC were carried out with *Chandra* and reported by Cseh et al. (2010). These authors found a number of sources located close to the core of the GC (sources #3, #7, and #12), with spectral properties similar to those of the X-ray binaries[4]. and averaged quiescent luminosities of $L_{\rm X}\simeq10^{32}$-10^{33} erg/s. The relatively small uncertainty in the position of IGR J17361-4441, as determined thanks to the *Chandra* observation of the source carried out by Pooley et al. (2011), suggests that the newly discovered *INTEGRAL* transient is associated with neither the objects detected previously in quiescence by Cseh et al. (2010), nor the COG of NGC6388 (see Fig. 5). The quiescent luminosity of IGR J17361-4441 can thus be estimated at $\sim10^{31}$ erg/s (Pooley et al. 2011), and is $\sim10^5$ times lower than it was at outburst. The radio observation discussed in Sect. 2.4, showed that the large increase in the X-ray luminosity from IGR J17361-4441 in outburst was not followed by any re-brightening in the radio domain. If we were to use our results obtained at 5.5 GHz and assume a distance to the source of 13.2 kpc, the rms level of 14.1 μJy derived in Sect. 2.4 from the ATCA data would translate into a 3σ upper limit on the radio luminosity from IGR J17361-4441 of $L_{\rm R}<5\times10^{28}$ erg/s (we adopted the same calculation procedure as in Cseh et al. 2010, see their Sect. 4). Assuming that the empirical relation between the mass of a BH and its radio and X-ray luminosity (the so-called "black-hole fundamental plane" Merloni et al. 2003; Körding et al. 2006) applies also in the present case, the measured values of $L_{\rm X}$ and $L_{\rm R}$ do not seem to be compatible with IGR J17361-4441 being a black-hole candidate (BHC) source, as the maximum allowed mass for this source would be far too low relative to expectations for these systems ($\gg M_{\odot}$).

As the broad-band spectral properties of the source discussed in Sect. 2.3 are reminiscent of what is observed for some X-ray binaries hosting accreting neutron stars (NSXRBs; see e.g., Psaltis 2004, for a review), we propose that an alternative possibility is that IGR J17361-4441 belongs to this class of objects. Highly magnetized ($\sim10^{12}$ G) NSs in high mass X-ray binaries (HMXBs) usually display in outburst X-ray spectra comprising a relatively flat absorbed power-law ($\Gamma\simeq0.5$-1.0) component plus a cut-off at higher energies (~10-50 keV, sometimes related to the strength of the NS magnetic field; Coburn et al. 2002). Soft excesses (<2 keV) are also observed from these sources in both quiescence and outburst, which are ascribed to either an accretion disk or a diffuse cloud around the compact object (see e.g., Hickox et al. 2004). Alternatively, a hot spot on the star can also give rise to a BB component in the spectrum with typical kT~0.3-1 keV. Given the results suggested by the XRT data analysis ($R_{\rm BB}\simeq1000$ km), the first case would probably be the most likely one for IGR J17361-4441. However, at odds with typical HMXBs in outburst, the new transient source did not display any pronounced variability in the X-ray emission on time-scales of neither a few hundreds of seconds (as in wind accreting systems) nor days (as in disk accreting systems). The relatively young age of typical HMXBs (few 10^6 yr) compared to that expected for the sources hosted in a globular cluster would also help us to exclude this hypothesis (the age of NGC6388 has been estimated to be around 10^{10} yr; Moretti et al. 2009). For this reason, IGR J17361-4441 might be more easily associated with a low mass X-ray binary (ages $\gtrsim10^8$ yr), even though these objects usually have softer spectra in the X-ray domain ($\Gamma\sim1.5$-2). The lack of any pulsation in the *RXTE* data could also not be used to firmly establish the presence of a NS in IGR J17361-4441. As suggested by Wijnands et al. (2011), a different possibility is that IGR J17361-4441 is a very faint X-ray transient (VFXT; Wijnands et al. 2006; Campana 2009). These objects have similarly hard spectra ($\Gamma\sim0.6$-1), and might emit a relatively stable X-ray flux in outburst even for a few years (see, e.g., the case of XMMU J174716.1281048; Del Santo et al. 2007). However, these sources typically have a lower X-ray luminosity (in the range 10^{34}-few$\times10^{35}$ erg/s) that that measured for IGR J17361-4441. Further observations of this new *INTEGRAL* transient with sensitive X-ray telescopes in the next few months might help us to understand the real nature of IGR J17361-4441.

Finally, we note that, as IGR J17361-4441 seems to be a new X-ray source in NGC6388 and not associated with the IMBH possibly residing at its center, the ATCA upper limit derived in Sect. 2.4 gives a significantly tighter constraint of the allowed mass for this object with respect to those reported previously. Following the discussion in Cseh et al. (2010), we assume that the quiescent X-ray luminosity of the IMBH is that determined by these authors with *Chandra* and consider the updated $L_{\rm R}$ provided by our ATCA ToO observation. Inserting these values into Eq.2 of Cseh et al. (2010) and taking into account the intrinsic scatter in the fundamental plane relation, the newly derived upper limit to the IMBH mass is $M<600\ M_{\odot}$. This is a factor of 2.5 smaller than that reported by Cseh et al. (2010).

Acknowledgements. We thank the Swift, RXTE and ATCA teams for the rapid scheduling of the ToO observations of IGR J17361-4441, and an anonymous referee for useful comments. TB acknowledges financial support from ASI (INAF-ASI I/009/10/0). AP acknowledges the support of the operating program of Regione Sardegna (ESF 2007-2013), L.R.7/2007, "Promotion of scientific research and technological innovation in Sardinia". This research has made use of the IGR Sources page maintained by J. Rodriguez & A. Bodaghee (http://irfu.cea.fr/Sap/IGR-Sources/).

References

Bozzo, E., Giunta, A., Stella, L., et al. 2009, A&A, 502, 21
Bradt, H. V., Rothschild, R. E., & Swank, J. H. 1993, Astronomy and Astrophysics Supplement Series (ISSN 0365-0138), 97, 355
Burrows, D. N., Hill, J. E., Nousek, J. A., et al. 2005, Space Sci. Rev., 120, 165
Campana, S. 2009, ApJ, 699, 1144
Coburn, W., Heindl, W. A., Rothschild, R. E., et al. 2002, ApJ, 580, 394
Courvoisier, T., Walter, R., Beckmann, V., et al. 2003, A&A, 411, L53
Cseh, D., Kaaret, P., Corbel, S., et al. 2010, MNRAS, 406, 1049
Dalessandro, E., Lanzoni, B., Ferraro, F. R., et al. 2008, ApJ, 677, 1069
Del Santo, M., Sidoli, L., Mereghetti, S., et al. 2007, A&A, 468, L17
Evans, P. A., Beardmore, A. P., Page, K. L., et al. 2009, MNRAS, 397, 1177
Ferrigno, C., Bozzo, E., Falanga, M., et al. 2011a, A&A, 525, A48+
Ferrigno, C., Bozzo, E., Rodriguez, J., & Gibaud, L. 2011b, The Astronomer's Telegram, 3566, 1
Gibaud, L., Bazzano, A., Bozzo, E., et al. 2011, The Astronomer's Telegram, 3565, 1
Heinke, C. O., Ruiter, A. J., Muno, M. P., & Belczynski, K. 2008, in American Institute of Physics Conference Series, Vol. 1010, A Population Explosion: The Nature & Evolution of X-ray Binaries in Diverse Environments, ed. R. M. Bandyopadhyay, S. Wachter, D. Gelino, & C. R. Gelino, 136–142
Hickox, R. C., Narayan, R., & Kallman, T. R. 2004, ApJ, 614, 881
Körding, E., Falcke, H., & Corbel, S. 2006, A&A, 456, 439
Lanzoni, B., Dalessandro, E., Ferraro, F. R., et al. 2007, ApJL, 668, L139
Lebrun, F., Leray, J. P., Lavocat, P., et al. 2003, A&A, 411, L141
Lund, N., Budtz-Jørgensen, C., Westergaard, N. J., et al. 2003, A&A, 411, L231
Merloni, A., Heinz, S., & di Matteo, T. 2003, MNRAS, 345, 1057
Moretti, A., Piotto, G., Arcidiacono, C., et al. 2009, A&A, 493, 539
Nucita, A. A., de Paolis, F., Ingrosso, G., Carpano, S., & Guainazzi, M. 2008, A&A, 478, 763
Pooley, D., Homan, J., Heinke, C. O., et al. 2011, The Astronomer's Telegram, 3627, 1
Psaltis, D. 2004, astro-ph/0410536
Ubertini, P., Lebrun, F., Di Cocco, G., et al. 2003, A&A, 411, L131
Vaughan, B. A., van der Klis, M., Wood, K. S., et al. 1994, ApJ, 435, 362
Wijnands, R., in't Zand, J. J. M., Rupen, M., et al. 2006, A&A, 449, 1117

---

[4] Source #3 appeared to be rather soft and possibly compatible with being a cataclysmic variable. We do not discuss here this hypothesis for IGR J17361-4441 as its X-ray luminosity seems to be far too high with respect to that typically observed from these objects (see e.g., Heinke et al. 2008).

Wijnands, R., Yang, Y., Degenaar, N., et al. 2011, The Astronomer's Telegram, 3595, 1
Wilson, W. E., Ferris, R. H., Axtens, P., et al. 2011, astro-ph/1105.3532
Winkler, C., Courvoisier, T., Di Cocco, G., et al. 2003, A&A, 411, L1